Think medicine safety when

Taking an OTC NSAID

What to know

NSAIDs (nonsteroidal anti-inflammatory drugs) are a commonly used type of pain reliever and fever reducer available over-the-counter (OTC) and by prescription.

The most well-known NSAIDs include ibuprofen, aspirin, and naproxen sodium.

NSAIDs are safe and effective when used as directed, but the risk of harmful side effects such as stomach bleeding, heart attack, or stroke increases the more you take and the longer you take them.

2 What you can do

Take the smallest effective dose and take it for the shortest amount of time needed.

NSAIDs*	AMOUNT	DOSE & FREQUENCY	DAILY LIMIT
Ibuprofen for example Motrin [®] IB and Advil [®]	200 mg per pill	1 pill every 4 to 6 hours while symptoms last (if pain or fever does not respond to 1 pill, 2 pills may be used)	1,200 mg 6 pills in 24 hours
Naproxen sodium for example Aleve*	220 mg per pill	1 pill every 8 to 12 hours while symptoms last (for the first dose, you may take 2 pills within the first hour)	660 mg 2 pills in any 8 to 12 hour period. Do not take more than 3 pills in 24 hours
Aspirin regular strength for example Bayer* Regular Strength	325 mg per pill	1 or 2 pills every 4 hours, or 3 pills every 6 hours	4,000 mg 12 pills in 24 hours
Aspirin extra strength for example	500 mg	1 or 2 pills every 4 to 6 hours	4,000 mg 8 pills in 24 hours



More than 900 OTC and Rx medicines contain an NSAID. Take ONLY 1 medicine that contains an NSAID at a time.



per pill

Write down all of the OTC and Rx medicines, vitamins, and herbal supplements you take. Bring the list and your bottles to your next visit.



If you have **any questions** about choosing or using OTC or Rx pain relievers, talk with your healthcare professional.

8 pills in 24 hours

Bayer® Extra Strength